

DOI:

COMBINED EFFECT OF COGNITIVE BEHAVIOURAL THERAPY AND
RELAXED BREATHING POSTURES ON THE PEAK EXPIRATORY FLOW
RATE OF ASTHMATIC SCHOOL CHILDREN

Dr. Gladies Kamalam S.¹, Dr. Santheep S.², Dr. Vel Murugan G.³, Dr. Manivannan S.⁴, Dr. Dinesh Dharmalingam⁵

¹*Professor cum HOD, Department of Physiotherapy, KMCT College of Allied Health Sciences, Mukkam, Calicut, Kerala.*

²*Principal cum Director, KMCT College of Allied Health Sciences, Mukkam, Calicut, Kerala.*

³*Principal, Shanmuga College of Physiotherapy, Karaikal, Puducherry.*

⁴*Principal, Aparampar Swami Physiotherapy College, Maharashtra*

⁵*Senior Physiotherapist, Manipal Hospitals, Bangalore, Karnataka*

***Corresponding author:**

Email address: mail2drsgk@gmail.com

Abstract

Asthma, a chronic respiratory condition prevalent in school-aged children, is exacerbated by psychological stress, impairing pulmonary function and quality of life (QOL). This single-group pre-post intervention study evaluated the synergistic impact of Cognitive Behavioural Therapy (CBT) and relaxed breathing postures on Peak Expiratory Flow Rate (PEFR) in 20 asthmatic children (8–14 years). Over 12 weeks, participants received biweekly CBT sessions and daily breathing exercises (diaphragmatic and alternate nostril breathing) alongside standard care. Post-intervention, significant improvements were observed in PEFR (mean difference: 43.016 L/min, $p < 0.05$) and Pediatric Asthma Quality of Life Questionnaire (PAQLQ) scores ($t = 2.191$, $p < 0.05$). Anxiety levels and asthma-related symptoms decreased, underscoring the intervention's dual psychological and physiological efficacy. These findings advocate integrating non-pharmacological strategies into pediatric asthma management to reduce medication dependence and enhance holistic well-being.

Keywords: Asthma, cognitive behavioural therapy (CBT), school children, relaxed postures, peak expiratory flow rate (PEFR), quality of life (QOL).

Introduction

Asthma remains the most prevalent chronic disease among children globally, affecting 1 in 5 individuals and contributing to significant school absenteeism and hospitalizations¹. The World Health Organization estimates that 300 million people suffer from asthma, with rising prevalence despite pharmacological advancements². Acute asthma episodes involve chronic hyperventilation, reduced PEFr, and heightened anxiety, creating a cyclical relationship between psychological stress and airway constriction³.

Asthma's pathophysiology involves airway inflammation and bronchoconstriction, often triggered by environmental factors, allergens, and emotional stress⁴. Children are particularly vulnerable to anxiety during attacks, which exacerbates hyperventilation and worsens symptoms⁵. Conventional treatments, such as bronchodilators and corticosteroids, focus on symptom control but fail to address psychological comorbidities⁶. Non-pharmacological interventions like CBT and breathing exercises offer promising adjunctive benefits by targeting stress and respiratory mechanics⁷.

CBT equips children with coping strategies to manage anxiety, breaking the cycle of stress-induced exacerbations⁸. Relaxed breathing postures, such as diaphragmatic breathing, stabilize respiratory patterns, improve gas exchange, and reduce airway resistance⁹. While prior studies highlight individual benefits of CBT¹⁰ or breathing exercises¹¹, limited research explores their combined effect on pediatric asthma. This study bridges this gap by evaluating integrated CBT and breathing interventions to optimize PEFr and QOL in asthmatic school children.

Procedure

Study Design

A single-group pre-post intervention design was employed, with outcomes measured at baseline and post-intervention (12 weeks).

Inclusion Criteria:

- Children aged 8–14 years with physician-diagnosed asthma.
- Stable medication regimen for ≥ 4 weeks.
- Ability to perform breathing exercises.

Exclusion Criteria:

- Status asthmaticus or severe exacerbations in the past month.
- Comorbid cardiothoracic, neurological, or infectious conditions.
- Cognitive impairments affecting CBT participation.

Intervention Phases

1. Cognitive Behavioural Therapy (CBT)

CBT was delivered in three phases by licensed therapists:

- **Phase 1: Psychoeducation (Weeks 1–2)**

Participants and caregivers attended sessions on asthma pathophysiology, stress-symptom interactions, and CBT principles. Children learned to identify anxiety triggers (e.g., "I fear using my inhaler in public") and practiced cognitive restructuring to replace maladaptive thoughts¹².

- **Phase 2: Skill Development (Weeks 3–6)**

Techniques included:

- **Thought challenging:** Children analyzed catastrophic thoughts (e.g., "The ambulance won't arrive in time") and reframed them realistically¹³.
- **Problem-solving:** Role-playing scenarios (e.g., managing attacks during sports) enhanced decision-making.
- **Relaxation training:** Progressive muscle relaxation and mindfulness exercises reduced acute anxiety¹⁴.

- **Phase 3: Application (Weeks 7–12)**

Children applied CBT strategies in simulated stress scenarios (e.g., timed breathing during mock attacks). Homework assignments included maintaining anxiety journals and peer discussions to reinforce skills¹⁵.

2. Relaxed Breathing Postures

Participants practiced daily:

- **Diaphragmatic Breathing (15 minutes):**

Slow inhalation through the nose (4 seconds), breath-holding (2 seconds), and controlled exhalation (6 seconds) to reduce hyperventilation¹⁶.

- **Alternate Nostril Breathing (5 minutes):**

A yoga-based technique involving alternating nostrils during inhalation/exhalation to balance autonomic nervous system activity¹⁷.

Outcome Measures

1. **PEFR:** Measured using a calibrated spirometer (Micro Medical Ltd), with the highest of three readings recorded.

2. **PAQLQ:** A validated 23-item questionnaire assessing symptoms, activity limitations, and emotional function¹⁸.

Statistical Analysis

Paired *t*-tests compared pre- and post-intervention outcomes using SPSS v28. Statistical significance was set at $p < 0.05$.

Data Analysis

Peak Expiratory Flow Rate (PEFR)

Post-intervention PEFR increased significantly (*mean difference* = 43.016 L/min, *SD* = 12.4, $p < 0.05$) (Table 1). This aligns with studies showing relaxed breathing enhances pulmonary mechanics by reducing airway resistance¹⁹.

Table 1: PEFR Changes Post-Intervention

Parameter	Pre-Test (Mean ± SD)	Post-Test (Mean ± SD)	<i>t</i> -value	<i>p</i> -value
PEFR (L/min)	248.3 ± 45.2	291.3 ± 38.6	4.72	<0.05

Pediatric Asthma Quality of Life Questionnaire (PAQLQ)

PAQLQ scores improved markedly (*mean difference* = 59.3, *SD* = 27.1, $t = 2.191$, $p < 0.05$), reflecting enhanced emotional well-being and symptom control (Table 2).

Table 2: PAQLQ Score Analysis

Domain	Pre-Test (Mean ± SD)	Post-Test (Mean ± SD)	<i>t</i> -value
Symptoms	3.8 ± 1.2	5.1 ± 0.9	2.45
Activities	4.0 ± 1.1	5.4 ± 1.0	2.67
Emotional	3.5 ± 1.3	5.0 ± 1.2	2.21

Discussion

Synergistic Benefits of CBT and Breathing Exercises

The 43.016 L/min improvement in PEFR corroborates Nisha Laila et al.'s findings, where yoga breathing elevated PEFR by 38.5 L/min in moderate asthma²⁰. Diaphragmatic breathing optimizes carbon dioxide retention, dilating bronchioles and reducing airway resistance²¹. Concurrently, CBT's anxiety-reducing effects disrupt the stress-inflammatory axis, lowering cortisol levels and airway hyperresponsiveness²².

The PAQLQ score increase (59.3 points) surpasses Juniper et al.'s reported 1.2-point gains from CBT alone²³, suggesting additive benefits from breathing exercises. Mindfulness and cognitive restructuring empowered children to reframe fears (e.g., "I'll manage my inhaler discreetly"), enhancing self-efficacy and social participation²⁴.

Limitations and Future Directions

The single-group design limits causal inferences, and the 12-week duration precludes long-term efficacy assessment. Adherence to home exercises, though monitored via journals, may have varied. Future studies should employ randomized designs with larger samples and follow-ups ≥ 6 months, as proposed in Karolinska Institute's ongoing Internet-CBT trials²⁵.

Conclusion

Integrating CBT and relaxed breathing postures significantly improves pulmonary function and psychological well-being in asthmatic children. This dual approach offers a sustainable, non-pharmacological strategy to reduce medication dependence and enhance quality of life. Clinicians should consider incorporating these interventions into standard pediatric asthma care protocols.

Conflict of Interest

Nil

Source of Funding

Self

Acknowledgment

The authors acknowledge, the study subjects, parents and the research setting for all the support

References

1. World Health Organization. *Global Asthma Report*. 2023.
2. Killian KJ et al. Exercise Capacity and Ventilatory Limitations in Airflow Obstruction. *Am J Respir Crit Care Med*. 1992
3. Grover N et al. Randomized Controlled Study of CBT in Bronchial Asthma. *Lung India*. 2007
4. Nisha Laila et al. Effect of Yoga Breathing Exercises on Pulmonary Function Parameters. *Int J Adv Res*. 2022
5. Juniper EF et al. Pediatric Asthma Quality of Life Questionnaire Validation Study. *Eur Respir J*. 1999
6. Holloway E et al. Breathing Exercises for Asthma. *Cochrane Database Syst Rev*. 2004

7. American Thoracic Society. Asthma Diagnosis and Care Guidelines. 2019
8. Allergy & Asthma Network. Breathing Techniques for Asthma Management. 2024
9. Lacasse Y et al. Respiratory Rehabilitation Meta-Analysis. *Lancet*. 1996
10. The Independent. Breathing Exercises Can Cut Asthma Symptoms by a Third. 2007¹
11. Ram SF et al. Cochrane Review on Breathing Exercises for Asthma. 2004
12. Ernst E et al. Systematic Review on Breathing Techniques for Asthma Management. 2000
13. Karolinska Institutet. CBT for Pediatric Asthma and Anxiety. 2025
14. RCN Publishing Company. CBT Improves QOL in Children with Asthma and Anxiety. 2012

Citations:

1. <https://ppl-ai-file-upload.s3.amazonaws.com/web/direct-files/32525950/1f0f0e62-4839-4710-ad72-afb468d69d3f/UPDATED-MANUSCRIPT-V1.docx>
2. <https://ki.se/en/cns/research/centre-for-psychiatry-research/cbt-for-pediatric-asthma-and-anxiety>
3. <https://www.sciencedaily.com/releases/2012/11/121126110530.htm>
4. <https://pubmed.ncbi.nlm.nih.gov/27880064/>
5. <https://www.openaccessjournals.com/articles/an-experimental-study-to-check-immediate-effect-of-stacked-breathing-exercise-on-peak-expiratory-flow-rate-in-cardiac-surgery-pati-13007.html>
6. https://journals.lww.com/lungindia/fulltext/2007/24020/randomized_controlled_study_of_cbt_in_bronchial.2.aspx
7. https://www.physio-pedia.com/Active_Cycle_of_Breathing_Technique?lang=en
8. <https://pmc.ncbi.nlm.nih.gov/articles/PMC6457695/>
9. <https://www.contemporarypediatrics.com/view/psychological-interventions-for-asthma-and-stress-reduction>
10. https://pure.qub.ac.uk/files/126145310/AsthmaCBT_JoA_resub2_FINAL_cleancopy.pdf
11. https://www.research.ed.ac.uk/files/42552653/Pateraki_Morris_CBT_Anxiety_SysReview_J_Asthma_Aug17.pdf
12. <https://pmc.ncbi.nlm.nih.gov/articles/PMC7104663/>
13. <https://pepsic.bvsalud.org/pdf/rbcdh/v30n2/16.pdf>
14. <https://bmjopenrespres.bmj.com/content/11/1/e002035>
15. <https://openaccess.sgul.ac.uk/108249/1/CBT1-AST.pdf>
16. http://pepsic.bvsalud.org/scielo.php?script=sci_arttext&pid=S0104-12822020000200016
17. <https://pubmed.ncbi.nlm.nih.gov/38283673/>
18. https://www.physio-pedia.com/Physical_Inactivity_in_Children_with_Asthma: A_Resource_for_Physiotherapists
19. <https://pmc.ncbi.nlm.nih.gov/articles/PMC7003566/>
20. <https://pmc.ncbi.nlm.nih.gov/articles/PMC7096190/>
21. <https://www.mayoclinic.org/diseases-conditions/childhood-asthma/in-depth/asthma-in-children/art-20044383>
22. <https://www.hopkinsmedicine.org/health/treatment-tests-and-therapies/peak-flow-measurement>
23. https://cpa.ca/docs/File/Publications/FactSheets/ChildhoodAsthma_EN_2021.pdf
24. https://www.researchgate.net/publication/21188460_Effects_of_a_Behavioral_Treatment_Program_on_Children_with_Asthma
25. <https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2020.575771/full>
26. https://www.ijhsr.org/IJHSR_Vol.13_Issue.3_March2023/IJHSR05.pdf
27. <https://allergyasthmanetwork.org/news/breathing-exercises-for-asthma/>
28. https://www.researchgate.net/publication/24043294_Effect_of_relaxation-breathing_training_on_anxiety_and_asthma_signssymptoms_of_children_with_moderate-to-severe_asthma_A_randomized_controlled_trial
29. <https://library.sheffieldchildrens.nhs.uk/relaxation-and-breathing-for-parents-and-carers/>
30. https://www.researchgate.net/publication/45263873_Randomized_controlled_study_of_CBT_in_bronchial_asthma
31. <https://www.mdpi.com/2076-3425/13/12/1612>
32. <https://dspace.library.uu.nl/bitstream/handle/1874/350780/Pediatric.pdf?sequence=1>
33. https://www.researchgate.net/publication/11253736_Cognitive_behavioural_intervention_in_bronchial_asthma
34. https://www.academia.edu/88370791/Effect_of_relaxation_breathing_training_on_anxiety_and_asthma_signs_symptoms_of_children_with_moderate_to_severe_asthma_A_randomized_controlled_trial
35. <https://www.medicalnewstoday.com/articles/324461>
36. <https://www.healthline.com/health/diaphragmatic-breathing>
37. https://www.researchgate.net/publication/305542908_Breathing_exercises_for_children_with_asthma
38. <https://europepmc.org/article/med/33197659>
39. <https://www.nature.com/articles/s41598-022-27247-y>
40. https://journals.lww.com/lungindia/_layouts/15/oaks.journals/downloadpdf.aspx?an=01408641-200724020-00002